

OR-25


TRANSITION METALL-FREE C-H/C-H AND C-LI/C-H COUPLING IN THE SYNTHESIS OF NEW AZAHETEROCYCLES DERIVATIVES

**Timofey D. Moseev,^a Mikhail V. Varaksin,^{a,b} Egor A. Nikiforov,^a Denis A. Gorlov^a
Oleg N. Chupakhin,^{a,b} Valery N. Charushin^{a,b}**

^aUral Federal University, 19 Mira St., 620002, Ekaterinburg, Russia

^bInstitute of Organic Synthesis, Ural Branch of the Russian Academy of Sciences,
22 S. Kovalevskaya St., 620041 Ekaterinburg, Russia. E-mail: chupakhin@ios.uran.ru

Abstract. Over the past decades, a progress in the synthesis via transition metal-free C–H bond functionalization reactions has been observed. This green chemistry-oriented strategy is highly attractive since it allows one to avoid protecting groups, to shorten the reaction time, to reduce. Reactions of nucleophilic substitution of hydrogen (S_N^H), being one of the promising mode of C–H functionalization methodology, has been used in our research project as a basic approach in two different synthetic schemes. The first one is direct C–H/C–H coupling of imidazole N-oxides with C-nucleophiles (pyrroles and phenols) by action of the activated starting substrate with acylating agent resulted in the substituted imidazoles as promising anti-radical and neuroprotection activity agents (Scheme 1 left). The second one is direct C–Li/CH coupling of pentafluorophenyl lithium, which was obtained *in situ* from pentafluorobenzene and *n*-BuLi, with azine or azole N-oxides was found to afford polyfluorinated azaheterocycles of special interest as new materials with photoluminescent properties (Scheme 1 right).



Scheme 1. C-H/C-H and C-Li/C-H coupling in the synthesis of new azaheterocycles derivatives.

As a result, transition metal-free C–H/C–H and C–Li/C–H coupling strategies were successfully applied to 2*H*-imidazole-N-oxides and pentafluorophenyllithium in the synthesis of novel azaheterocycles derivatives of the particular interest in medicinal and coordinational chemistry, photoactive materials etc.

References

1. Metal-free C–H functionalization of 2*H*-imidazole 1-oxides with pyrrolyl fragments in the design of novel azaheterocyclic ensembles / M.V. Varaksin, T.D. Moseev, O.N. Chupakhin, V.N. Charushin [et al.] // Org. Biomol. Chem. – 2017. – Vol. 15. – P. 8280-8284.
2. Direct C–Li/C–H coupling of pentafluorophenyl lithium with azines - An atom- and step-economical strategy for the synthesis of polyfluoroaryl azaaromatics / M.V. Varaksin, T.D. Moseev, V.N. Charushin [et al.] // J. Organomet. Chem. – 2018. – Vol. 867. – P. 278-283.

The study was carried out with the financial support of the Russian Foundation for Basic Research in the framework of a research project 18-33-00226 and Russian Science Foundation as part of a research project 18-73-00088.